



Learning if HARVONI is right for you starts with speaking to a Hep C Specialist. HARVONI is a prescription medicine used to treat adults with chronic (lasting a long time) hepatitis C (Hep C) genotype (GT) 1, 4, 5 or 6 infection with or without cirrhosis (compensated). In those with GT 1 and advanced cirrhosis (decompensated) or with GT 1 or 4 with or without cirrhosis (compensated) who have had a liver transplant, HARVONI is used with ribavirin.

Use this guide to help your Hep C Specialist understand you're ready to put Hep C behind you.



Be prepared to tell your Hep C Specialist about your medical history, including your history with Hep C and any previous Hep C treatment. Hep C experience is one of the factors that helps your specialist determine the length of your treatment.

□ 0-10 years □ 11-20 years □ 21+ years □ Not sure B. Have you previously been treated for Hep C? □ Yes □ No	
	_
☐ Yes ☐ No	
C. Do you have cirrhosis?	-
☐ Yes ☐ No ☐ Not sure	
D. List anything else that may be affecting your health right now, such as hepatitis B infectio (or if you have ever had hepatitis B infection), liver problems other than Hep C or if you had a liver transplant; kidney problems or if you are on dialysis; HIV infection; or any othe medical condition.	have

Important Safety Information

What is the most important information I should know about HARVONI?

HARVONI can cause serious side effects, including:

• **Hepatitis B virus reactivation:** Before starting HARVONI treatment, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during and after treatment with HARVONI. This may cause serious liver problems, including liver failure and death. If you are at risk, your healthcare provider will monitor you during and after taking HARVONI.







Be prepared to tell your Hep C Specialist about symptoms you may be experiencing.

Let your specialist know about any of the symptoms below, or any other symptoms you may be experiencing—even if you don't think they seem related to your liver or hepatitis C.

	Which of the following symptoms have you had?				
	☐ Fever	Tiredness	☐ Loss of appetite	Nausea	
	☐ Vomiting	☐ Dark urine	☐ Abdominal pain	☐ Joint pain	
	☐ Gray-colored stools	☐ Gray-colored stools ☐ Jaundice (yellowing of skin or eyes)			
	Other				
3	Be prepared to tell your Hep C Specialist about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. HARVONI may affect the way other medicines work, and other medicines may affect how HARVONI works or cause side effects.				
	List all the medicines (prescription and over-the-counter), vitamins, and herbal supplements you take.				

Important Safety Information (continued)

What should I tell my healthcare provider before taking HARVONI?

- Tell your healthcare provider about all of your medical conditions, including if you have ever had hepatitis B infection, liver problems other than hepatitis C infection, or a liver transplant; if you have severe kidney problems or are on dialysis; if you have HIV; or if you are pregnant or breastfeeding or plan to become pregnant or breastfeed. It is not known if HARVONI will harm your unborn baby or pass into your breast milk. If you take HARVONI with ribavirin, you should also read the ribavirin Medication Guide for important pregnancy-related information.
- Tell your healthcare provider and pharmacist about all the medicines you take, including prescription
 and over-the-counter medicines, vitamins, and herbal supplements. HARVONI and certain other
 medicines may affect each other, or may cause side effects.







Express yourself and be clear. Remember, you are your best advocate. Explain how Hep C makes you feel. Provide as much specific information as possible.

How exactly does Hep C make you feel? How have things changed over time?				

Important Safety Information (continued)

What are the possible side effects of HARVONI?

Serious side effects may also include:

• Slow heart rate (bradycardia): HARVONI, when taken with amiodarone (Cordarone®, Nexterone®, Pacerone®), a medicine used to treat certain heart problems, may cause slow heart rate, which in some cases has led to death or the need for a pacemaker. Get medical help right away if you take amiodarone with HARVONI and get any of the following symptoms: fainting or near-fainting, dizziness or lightheadedness, not feeling well, weakness, extreme tiredness, shortness of breath, chest pains, confusion, or memory problems.

The most common side effects of HARVONI include tiredness, headache and weakness.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Important Facts about HARVONI including Important Warning on hepatitis B reactivation located on page 4.



(har-VOE-nee)

MOST IMPORTANT INFORMATION ABOUT HARVONI

HARVONI can cause serious side effects, including:

Hepatitis B virus reactivation: Before starting treatment with HARVONI, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during or after treatment with HARVONI. This may cause serious liver problems, including liver failure and death. Your healthcare provider will monitor you if you are at risk for hepatitis B reactivation during and after you stop taking HARVONI.

ABOUT HARVONI

HARVONI is a prescription medicine used to treat people 12 years of age and older (or weighing at least 77 pounds) with chronic (lasting a long time) hepatitis C genotype (GT) 1, 4, 5 or 6 infection with or without cirrhosis (compensated). In adults with GT 1 infection and advanced cirrhosis (decompensated) or with GT 1 or 4 infection with or without cirrhosis (compensated) who have had a liver transplant, HARVONI is used with ribayirin.

POSSIBLE SIDE EFFECTS OF HARVONI

HARVONI can cause serious side effects, including:

- Those in the "Most Important Information about HARVONI" section.
- Slow heart rate (bradycardia): HARVONI, when taken with amiodarone (Cordarone®, Nexterone®, Pacerone®), a medicine used to treat certain heart problems, may cause slow heart rate, which in some cases has led to death or the need for a pacemaker. Get medical help right away if you take amiodarone with HARVONI and get any of the following symptoms:
- fainting or near-fainting
- dizziness or lightheadedness
- not feeling well
- weakness
- extreme tiredness
- shortness of breath
- chest pains
- confusion
- memory problems

The most common side effects of HARVONI include tiredness, headache and weakness.

These are not all the possible side effects of HARVONI. Tell your healthcare provider if you have any new symptoms while taking HARVONI.

IMPORTANT FACTS

This is only a brief summary of important information about HARVONI and does not replace talking to your healthcare provider about your condition and your treatment.

BEFORE TAKING HARVONI

Tell your healthcare provider about all of your medical conditions, including if you:

- have ever had hepatitis B virus infection
- have liver problems other than hepatitis C infection
- had a liver transplant
- have severe kidney problems or are on dialysis
- have HIV infection
- are pregnant or breastfeeding, or plan to become pregnant or breastfeed
 - If you take HARVONI with ribavirin, you should also read the ribavirin Medication Guide for important pregnancy-related information.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider.
- HARVONI and certain medicines may affect each other, or cause side effects.

GET MORE INFORMATION

- This is only a brief summary of important information about HARVONI. Talk to your healthcare provider or pharmacist to learn more.
- Go to HARVONI.com or call 1-844-READY41.
- To learn about potential savings, call 1-855-7-MYPATH or go to HARVONI.com/support

